

ABOUT THE DIAGNOSIS

Uveitis is inflammation within the eye. The structures affected are the iris (the colored part of the front of the eye) and the blood vessel-rich layer that lies deep inside the eye, beneath the retina, called the choroid. Uveitis is a common condition in both cats and dogs and has many causes. Untreated uveitis can seriously damage the eye and even cause blindness.

Eye disorders that may cause uveitis include tumors, cataracts, severe corneal ulcers, high blood pressure, immune-mediated conditions, some infections, and trauma such as being hit by a car. In both dogs and cats, several fungal diseases are known to cause uveitis. In cats, feline leukemia virus infection, feline immunodeficiency virus infection, and feline infectious peritonitis can all be associated with uveitis. A great variety of other infectious diseases can have uveitis as one of the signs. Cats may develop a form of chronic uveitis where no underlying cause can be found. In summary, once uveitis is identified by a veterinarian, many potential causes are possible and some tests need to be performed in order to pinpoint the underlying cause and treat it.

Pets with uveitis may have variety of signs. The most noticeable to pet owners are redness of the eye, squinting, tearing, and avoidance of bright light. The eye may be painful. The clear part of the eye may become hazy or cloudy or, in severe cases, contain white flecks (precipitates). Eventually blindness can occur.

Many eye disorders appear like uveitis, and uveitis occurs as a secondary condition in several eye and body-wide diseases. Your veterinarian will conduct a complete eye exam, including using an ophthalmoscope to evaluate the back of the eye. If uveitis or glaucoma is suspected, the intraocular pressure will be measured using an instrument called a tonometer. In early uveitis, the intraocular pressure is lower than normal, while cases that have progressed may develop glaucoma where the pressure is abnormally high. Pressures and other tests help define the uveitis in an individual dog or cat and give a clearer idea of treatment choices and outlook for improvement (prognosis).

Unless the cause of your pet's uveitis is obvious, such as trauma, intraocular tumor, or cataract, your veterinarian will rely on additional diagnostic tests to find the underlying disease. Blood tests can be used for detecting many infectious diseases. Chest or abdominal x-rays may be required to look for tumors or signs of fungal diseases. Ultrasound examination of the eye may be recommended in cases when the eye is too cloudy for a thorough examination of the eye with an ophthalmoscope or to check for traumatic rupture of the eyeball or foreign bodies.

LIVING WITH THE DIAGNOSIS

The long-term outlook (prognosis) for uveitis is variable, depending on the underlying cause. Some of the body-wide infections that are associated with uveitis carry a guarded prognosis; they can become

life-threatening if the affected dog or cat does not improve with treatment or may improve significantly and resolve completely. In the case of uveitis caused by trauma or another treatable disease, the chance of resolution without serious damage to the eye depends upon the severity of the inflammation; some patients recover very well and regain normal vision and comfort, while others are left with some degree of complications such as blindness, cataracts, glaucoma, or a shrunken eye.

TREATMENT

Prompt and intensive treatment of uveitis is necessary to prevent damage to the eye. The underlying disease causing the uveitis must be controlled, if possible. Severe cases may require in-hospital treatment initially. Antiinflammatory drugs are used to control the inflammation in the eye. Corticosteroids (cortisone-like drugs) are the most frequently used medications. Depending on the underlying cause and the severity of the condition, these may be given as eye drops or eye ointments (applied to the eye several times per day). In some cases, corticosteroids or immunosuppressive drugs may be given as tablets by mouth. To reduce pain and prevent some other changes in the eye caused by the uveitis, atropine eye drops or ointment may be prescribed. Treatment of uveitis typically needs to continue for at least 2 months since the eye tissues often are slow to return to normal.

DOs

- When using eye drops or ointments to treat your pet, be sure to thoroughly wash your hands after treatment. This is especially important with atropine medications, which will cause your pupils to dilate if you accidentally get them in your eyes, and antibiotics, which can cause severe allergic reactions in susceptible people.
- Keep track of your dog's or cat's symptoms and vision. Day-to-day fluctuations are common, and it is the trend over several days to weeks that determines the best course of treatment.
- Consider a referral to a veterinary ophthalmologist (www.acvo.org) for the expert evaluation of a veterinary eye specialist.

DON'Ts

- Do not discontinue treatment before advised to do so by your veterinarian. Even if the eye looks better, treatment is needed for several weeks to completely control the condition.

WHEN TO CALL YOUR VETERINARIAN

- If you are unable to apply eye medications as directed.
- If there are changes in the appearance of the eye.

SIGNS TO WATCH FOR

- Atropine may cause salivation or vomiting in some cats. Contact your veterinarian if this occurs.

ROUTINE FOLLOW-UP

- Initially: within a few days of the confirmation of uveitis to evaluate response to treatment. If response is good, further visits should be scheduled every 2 to 3 weeks to monitor progress.

Other information that may be useful: "How-To" Client Education Sheet:

- How to Administer Eye Medications

Practice Stamp or Name & Address

Also available in Spanish.